

G to GG QM Changes Compare Table

<p>Current Measure: Percent of Residents Who Made Improvements in Function (SS) (CMS ID: N037.03)</p>	<p>GG Equivalent Measure: Discharge Function Score (CMS ID: S042.01)</p>
<p>Measure Description: This measure reports the percentage of short-stay residents who were discharged from the nursing home that gained more independence in transfer, locomotion, and walking during their episodes of care.</p>	<p>Measure Description: This measure estimates the percentage of Medicare Part A SNF stays that meet or exceed an expected discharge function score.</p>
<p><u>Numerator</u> Short-stay residents who:</p> <ol style="list-style-type: none"> Have a change in performance score that is negative ([valid discharge assessment] - [valid preceding PPS 5-Day assessment or OBRA Admission assessment] < [0]). <p>Performance is calculated as the sum of G0110B1 (transfer: self-performance), G0110E1 (locomotion on unit: self-performance), and G0110D1 (walk in corridor: self-performance), with 7's (activity occurred only once or twice) and 8's (activity did not occur) recoded to 4's (total dependence).</p> <p><u>Denominator</u> Short-stay residents who: Meet all of the following conditions, except those with exclusions:</p> <ol style="list-style-type: none"> Have a valid discharge assessment (A0310F = [10]), and Have a valid preceding PPS 5-Day assessment (A0310B = [01]) or OBRA Admission assessment (A0310A = [01]). 	<p><u>Numerator</u> The total number of Medicare Part A SNF stays (Type 1 SNF Stays only) in the denominator, except those that meet the exclusion criteria, with an observed discharge function score that is equal to or greater than the calculated expected discharge function score.</p> <p><u>Denominator</u> The total number of Medicare Part A SNF stays (Type 1 SNF Stays only), except those that meet the exclusion criteria.</p> <p><u>GG Function items used for discharge function score calculations:</u></p> <ul style="list-style-type: none"> • GG0130A3. Eating • GG0130B3. Oral hygiene • GG0130C3. Toileting hygiene • GG0170A3. Roll left and right • GG0170C3. Lying to sitting on side of bed • GG0170D3. Sit to stand • GG0170E3. Chair/bed-to-chair transfer • GG0170F3. Toilet transfer • GG0170I3: Walk 10 Feet* • GG0170J3: Walk 50 Feet with 2 Turns* • GG0170R3. Wheel 50 feet with 2 Turns*
<p><u>Exclusions</u> Residents satisfying any of the following conditions:</p> <ol style="list-style-type: none"> 1.1. Comatose (B0100 = [1]) on the PPS 5-Day assessment or OBRA Admission assessment, whichever was used in the QM. 1.2 Life expectancy of less than 6 months (J1400 = [1]) on the PPS 5-Day assessment or OBRA Admission assessment, whichever was used in the QM. 	<p><u>Exclusions</u> Medicare Part A SNF stays are excluded if:</p> <ol style="list-style-type: none"> 1. The Medicare Part A SNF stay is an incomplete stay: Unplanned discharge - Discharge to acute hospital, psychiatric hospital, long-term care hospital indicated - SNF PPS Part A stay less than 3 days - The resident died during the SNF stay 2. The resident has any of the following medical conditions at the time of admission (i.e., on the 5-Day PPS assessment): Coma,

G to GG QM Changes Compare Table

<p>1.3 Hospice (O0100K2 = [1]) on the PPS 5-Day assessment or OBRA Admission assessment, whichever was used in the QM.</p> <p>1.4 Information on Transfer: self-performance, walk in corridor: self-performance, or locomotion on unit: self-performance is <u>missing on any of the assessments used to calculate the QM</u> (G0110B1, G0110D1, or G0110E1 = [-]) (i.e., valid discharge assessment, and PPS 5-Day assessment or OBRA Admission assessment, whichever was used in the QM).</p> <p>1.5 Residents with no impairment (sum of G0110B1, G0110D1 and G0110E1 = [0]) on the PPS 5-Day assessment or OBRA Admission assessment, whichever was used in the QM.</p> <p>1.6. Residents with an unplanned discharge on any assessment during the care episode (A0310G = [2])</p>	<p>persistent vegetative state, complete tetraplegia, severe brain damage, locked-in syndrome, or severe anoxic brain damage, cerebral edema or compression of brain.</p> <p>3. The resident is younger than age 18:</p> <p>4. The resident is discharged to hospice or received hospice while a resident:</p> <p>5. The resident did not receive physical or occupational therapy services at the time of admission (i.e., on the 5- Day PPS assessment):</p>
<p><u>Covariates</u></p> <ol style="list-style-type: none"> Age on the PPS 5-Day assessment Gender Severe cognitive impairment Long Form ADL (LFADL) Scale (G0110A1 + G0110B1 + G0110E1 + G0110G1 + G0110H1 + G0110I1 + G0110J1). If any (G0110A1, G0110B1, G0110E1, G0110G1, G0110H1, G0110I1, G0110J1) = [7, 8], recode the item to equal [4]. 	<p><u>Covariates</u></p> <ol style="list-style-type: none"> Age group Admission function – continuous form Admission function – squared form Primary medical condition category Interaction between admission function and primary medical condition category Prior surgery Prior functioning: self-care Prior functioning: indoor mobility (ambulation) Prior functioning: stairs Prior functioning: functional cognition Prior mobility device use Stage 2 pressure ulcer/injury Stage 3, 4, or unstageable pressure ulcer/injury Cognitive abilities Communication impairment Urinary Continence Bowel Continence History of falls Nutritional approaches High BMI Low BMI Comorbidities

G to GG QM Changes Compare Table

<p style="text-align: center;">Current Measure: Percent of Residents Whose Ability to Move Independently Worsened (LS) (CMS ID: N035.03)</p>	<p style="text-align: center;">GG Equivalent Measure: Percent of Residents Whose Ability to Walk Independently Worsened (LS) (CMS ID: N035.04)</p>
<p>Measure Description: This measure reports the percent of long-stay residents who experienced a decline in independence of locomotion during the target period.</p>	<p>Measure Description: This measure reports the percent of long-stay residents who experienced a decline in independence of locomotion during the target period.</p>
<p><u>Numerator</u> Long-stay residents with a selected target assessment and at least one qualifying prior assessment who have a decline in locomotion when comparing their target assessment with the prior assessment. Decline identified by:</p> <ol style="list-style-type: none"> 1. Recoding all values (G0110E1 = [7, 8]) to (G0110E1 = [4]). 2. An increase of one or more points on the "locomotion on unit: self-performance" item between the target assessment and prior assessment (G0110E1 on target assessment – G0110E1 on prior assessment ≥1) <p><u>Denominator</u> Long-stay residents who have a qualifying MDS 3.0 target assessment and at least one qualifying prior assessment, except those with exclusions.</p>	<p><u>Numerator</u> Long-stay residents with a selected target assessment and at least one qualifying prior assessment who have a decline in locomotion when comparing their target assessment with the prior assessment. Decline identified by:</p> <ol style="list-style-type: none"> 1. Recoding all values (GG0170I = [07, 09, 10, 88]) to (GG0170I = [01]). 2. A decrease of one or more points on the "Walk 10 feet" item between the target assessment and prior assessment (GG0170I on target assessment – GG0170I on prior assessment ≤ -1) <p><u>Denominator</u> Long-stay residents who have a qualifying target assessment and at least one qualifying prior assessment, except those with exclusions.</p>
<p><u>Exclusions (Any of the following)</u></p> <ol style="list-style-type: none"> 1. Comatose or missing data on comatose at the prior assessment. 2. Prognosis of less than 6 months at the prior assessment as 3. Resident totally dependent during locomotion on prior assessment 4. Missing data on locomotion on target or prior assessment 5. Prior assessment is a discharge with or without return anticipated 6. No prior assessment is available to assess prior function. 	<p><u>Exclusions (Any of the following)</u></p> <ol style="list-style-type: none"> 1. Comatose or missing data on comatose at the prior assessment. 2. Prognosis of less than 6 months at the prior assessment 3. Resident dependent or activity was not attempted during locomotion on prior assessment 4. Missing data on locomotion on target or prior assessment 5. Prior assessment is a discharge with or without return anticipated 6. No prior assessment is available to assess prior function. 7. Prior or target assessment dates before 10/01/2023
<p><u>Covariates</u></p> <ol style="list-style-type: none"> 1. Eating (self-performance) from prior assessment 2. Toileting (self-performance) from prior assessment 3. Transfer (self-performance) from prior assessment 4. Walking in Corridor (self-performance) from prior assessment 5. Severe cognitive impairment from prior assessment 	<p><u>Covariates</u></p> <ol style="list-style-type: none"> 1. Eating from prior assessment 2. Toilet Transfer from prior assessment 3. Sit to Stand from prior assessment 4. Walk 10 Feet from prior assessment 5. Severe cognitive impairment from prior assessment

G to GG QM Changes Compare Table

<p>6. Linear Age 7. Gender 8. Vision 9. Oxygen use 10. All covariates are missing if no prior assessment is available.</p>	<p>6. Linear Age 7. Gender. 8. Vision. 9. Oxygen use. 10. All covariates are missing if no prior assessment is available.</p>
--	---

G to GG QM Changes Compare Table

<p>Current Measure: Percent of Residents Whose Need for Help with Activities of Daily Living Has Increased (LS) (CMS ID: N028.02)</p>	<p>GG Equivalent Measure: Percent of Residents Whose Need for Help with Activities of Daily Living Has Increased (LS) (CMS ID: N028.03)</p>
<p>Measure Description: This measure reports the percent of long-stay residents whose need for help with late-loss Activities of Daily Living (ADLs) has increased when compared to the prior assessment.</p>	<p>Measure Description: This measure reports the percent of long-stay residents whose need for help with late-loss Activities of Daily Living (ADLs) has increased when compared to the prior assessment.</p>
<p><u>Numerator</u> Long-stay residents with selected target and prior assessments that indicate the need for help with late-loss Activities of Daily Living (ADLs) has increased when the selected assessments are compared.</p> <p>The four late-loss ADL items are self-performance bed mobility (G0110A1), self-performance transfer (G0110B1), self-performance eating (G0110H1), and self-performance toileting (G0110I1).</p> <p>An increase is defined as: - an increase in two or more coding points in one late-loss ADL item or - one point increase in coding points in two or more late-loss ADL items.</p> <p>Note that for each of these four ADL items, if the value is equal to [7, 8] on either the target or prior assessment, then recode the item to equal [4] to allow appropriate comparison.</p> <p><u>Denominator</u> All long-stay residents with a selected target and prior assessment, except those with exclusions.</p>	<p><u>Numerator</u> Long-stay residents with selected target and prior assessments that indicate the need for help with late-loss Activities of Daily Living (ADLs) has increased when the selected assessments are compared.</p> <p>The four late-loss ADL items are Sit to Lying (GG0170B), Sit to Stand (GG0170D), Eating (GG0130A), and Toilet Transfer (GG0170F).</p> <p>An increase in need for help is defined as: - a decrease in two or more coding points in one late-loss ADL item or - one point decrease in coding points in two or more late-loss ADL items.</p> <p>Note that for each of these four ADL items, if the value is equal to [07, 09, 10, 88] on either the target or prior assessment, then recode the item to equal [01] to allow appropriate comparison.</p> <p><u>Denominator</u> All long-stay residents with a selected target and prior assessment, except those with exclusions.</p>
<p><u>Exclusions</u></p> <ol style="list-style-type: none"> 1. All four of the late-loss ADL items indicate total dependence, activity occurred only once or twice, or activity did not occur on the prior assessment 2. Three of the late-loss ADLs indicate total dependence on the prior assessment, as in #1 AND the fourth late-loss ADL indicates extensive assistance on the prior assessment. 	<p><u>Exclusions</u></p> <ol style="list-style-type: none"> 1. All four of the late-loss ADL items indicate dependence or activity was not attempted on the prior assessment 2. Three of the late-loss ADLs indicate dependence or activity was not attempted on the prior assessment, as in exclusion 1 AND the fourth late-loss ADL indicates substantial/maximal assistance on the prior assessment.

G to GG QM Changes Compare Table

<p>3. If resident is comatose missing data on comatose on the target assessment.</p> <p>4. Prognosis of life expectancy is less than 6 months on the target assessment.</p> <p>5. Hospice care on the target assessment.</p> <p>6. The resident is not in the numerator and data is missing for the four late loss ADLs on the prior or target assessment.</p>	<p>3. Comatose or missing data on comatose (B0100 = [1, -]) on the target assessment.</p> <p>4. Prognosis of life expectancy is less than 6 months on the target assessment.</p> <p>5. Hospice care on the target assessment.</p> <p>6. The resident is not in the numerator and data is missing for the four late loss ADLs on the prior or target assessment.</p> <p>7. No prior assessment is available to assess prior function.</p> <p>8. Prior or target assessment date before 10/01/2023.18</p>
<p><u>Covariates:</u> Not Applicable</p>	<p><u>Covariates:</u> Not Applicable</p>

G to GG QM Changes Compare Table

Current Measure: Percent of High-Risk Residents With Pressure Ulcers (LS)24 (CMS ID: N015.03)	GG Equivalent Measure: Percent of Residents With Pressure Ulcers (LS) (CMS ID: N045.01)
Measure Description: This measure captures the percentage of long-stay, high-risk residents with Stage II-IV or unstageable pressure ulcers	Measure Description: This measure captures the percentage of long-stay residents with Stage II-IV or unstageable pressure ulcers.
<p><u>Numerator</u> All long-stay residents with a selected target assessment that indicates Stage II-IV or unstageable pressure ulcers are present.</p> <p><u>Denominator</u> All long-stay residents with a selected target assessment that meet the definition of high risk, except those with exclusions.</p> <p>Residents are defined as high-risk if they meet one or more of the following three criteria on the target assessment:</p> <ol style="list-style-type: none"> 1. Impaired bed mobility or transfer 2. Comatose 3. Malnutrition or at risk of malnutrition 	<p><u>Numerator</u> All long-stay residents with a selected target assessment that indicates Stage II-IV or unstageable pressure ulcers are present.</p> <p><u>Denominator</u> All long-stay residents with a selected target assessment except those with exclusions.</p>
<p><u>Exclusions</u></p> <ol style="list-style-type: none"> 1. Target assessment is an OBRA Admission assessment (A0310A = [01]) or a PPS 5-Day assessment (A0310B = [01]) 2. If the resident is not included in the numerator (the resident did not meet the pressure ulcer conditions for the numerator) and no data is available for the Stage II-IV or unstageable pressure ulcers items on the target assessment 	<p><u>Exclusions</u></p> <ol style="list-style-type: none"> 1. Target assessment is an ORBA Admission assessment (A0310A = [01]) or a PPS 5-Day assessment (A0310B = [01]) 2. If the resident is not included in the numerator and no data is available for the Stage II-IV or unstageable pressure ulcers items on the target assessment 3. Assessments with target dates before 10/01/2023.
<p><u>Covariates:</u> Not Applicable</p>	<p><u>Covariates</u></p> <ol style="list-style-type: none"> 1. Impaired Functional Mobility: Lying to Sitting on Side of Bed on 2. Bowel Incontinence on target assessment. 3. Diabetes Mellitus, Peripheral Vascular Disease or Peripheral Arterial Disease on target assessment. 2. Indicator of low body mass index based on height (K0200A) and weight (K0200B) on target assessment. 3. Malnutrition or at risk of malnutrition on target assessment. 4. Dehydrated on target assessment. 5. Infections: Septicemia, Pneumonia, Urinary Tract Infection or Multidrug-Resistant Organism on target assessment. 6. Moisture Associated Skin Damage on target assessment.



"A *Knowledgeable* and *Compassionate* partner"

G to GG QM Changes Compare Table

	7. Hospice Care on target assessment.
--	---------------------------------------

G to GG QM Changes Compare Table

<p>Current Measure: Percent of Low-Risk Residents Who Lose Control of Their Bowel or Bladder (LS) (CMS ID: N025.02)</p>	<p>GG Equivalent Measure: Percent of Residents With New or Worsened Bowel or Bladder Incontinence (LS) (CMS ID: N046.01)</p>
<p>Measure Description: The measure reports the percent of long-stay residents who frequently lose control of their bowel or bladder.</p>	<p>Measure Description: This measure reports the percent of long-stay residents with new or worsened bowel or bladder incontinence between the prior assessment and target assessment.</p>
<p><u>Numerator</u> Long-stay residents with a selected target assessment that indicates frequently or always incontinence of the bladder</p> <p><u>Denominator</u> All long-stay residents with a selected target assessment, except those with exclusions.</p>	<p><u>Numerator</u> Long-stay residents with selected target and prior assessments that indicate a new or worsened case of bowel or bladder incontinence has occurred when the selected assessments are compared.</p> <p>Residents meet the definition of new or worsened bowel or bladder incontinence if any of the following conditions are true:</p> <p>Condition A: A new case of bowel incontinence is defined as an increase in one or more coding points on the bowel continence item (H0400) from always continent to either occasionally, frequently, or always incontinent.</p> <p>Condition B: A worsened case of bowel incontinence is defined as an increase in one or two coding points on the bowel continence item (H0400) from occasionally incontinent to frequently or always incontinent or from frequently incontinent to always incontinent.</p> <p>Condition C: A new case of bladder incontinence is defined as an increase in one or more coding points on the bladder continence item from always continent or occasionally incontinent to frequently or always incontinent.</p> <p>Condition D: A worsened case of bladder incontinence is defined as an increase in one coding point on the bladder continence item (H0300) from frequently incontinent to always incontinent.</p>

G to GG QM Changes Compare Table

	<p><u>Denominator</u> All long-stay residents with a selected target and prior assessment, except those with exclusions.</p>
<p><u>Exclusions</u></p> <ol style="list-style-type: none"> 1. Target assessment is an admission assessment or a PPS 5-Day assessment 2. Resident is not in numerator and data is missing for the incontinence items. 3. Residents who have any of the following high-risk conditions: <ol style="list-style-type: none"> 3.1. Severe cognitive impairment on the target assessment 3.2. Totally dependent in bed mobility self-performance 3.3. Totally dependent in transfer self-performance 3.4. Totally dependent in locomotion on unit self-performance 4. Resident does not qualify as high risk and both of the following two conditions are true for the target assessment: <ol style="list-style-type: none"> 4.1. BIMS summary score is missing and 4.2. Short term memory data is missing 5. Resident does not qualify as high risk and any of the following three conditions are true: <ol style="list-style-type: none"> 5.1. Data for bed mobility is missing 5.2. Data for transfers is missing 5.3. Data for locomotion on unit is missing 6. Resident is comatose or comatose status is missing on the target assessment. 7. Resident has an indwelling catheter or indwelling catheter status is missing on the target assessment. 8. Resident has an ostomy or ostomy status is missing on the target assessment. 	<p><u>Exclusions</u></p> <ol style="list-style-type: none"> 1. Target assessment is an admission assessment or a PPS 5-Day assessment 2. Resident is not in numerator and data is missing for the incontinence items on the prior assessment or on the target assessment. 3. Resident is comatose or comatose status is missing on the prior assessment, or on the target assessment. 4. Resident has an indwelling catheter or indwelling catheter status is missing on the prior assessment, or on the target assessment. 5. Resident has an ostomy or ostomy status is missing on the prior assessment, or on the target assessment. 6. No prior assessment is available to assess prior function. 7. Prior or target assessments with dates before 10/01/2023.
<p><u>Covariates</u> Not Applicable</p>	<p><u>Covariates</u></p> <ol style="list-style-type: none"> 1. Severe cognitive impairment on target assessment. 2. Sit to Lying on prior assessment. 3. Sit to Stand on prior assessment.